



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

x, of the equations (14), are precisely the only possible solutions of the given problem of the Calculus of Variations; or, the only possible solutions of the given problem of the Calculus of Variations are given by the characteristic torsion strips of the equation (2), regarded as a differential equation.

We might now easily go on to set up, as we have done in Part I, § 6 for equations of the first order, the equations of the common characteristics of two partial differential equations of the second order, and the condition that they be in involution, regarding the common characteristics as the common possible solutions of two problems in the Calculus of Variations, which reduce to a single problem of the type given above with one additional auxiliary condition of the type (2). Finally, we might seek the characteristics for two differential equations of the first order in two dependent variables, and so on. These problems will, however, offer no essential difficulty to the reader, and we will not enter into a discussion of them here; the main point of the existence of a connection between the theory of characteristics and the Calculus of Variations already having been demonstrated.

SHEFFIELD SCIENTIFIC SCHOOL,
YALE UNIVERSITY.
FEBRUARY, 1903.

ERRATUM.

Page 139, line 7: instead of $q = \psi'(y)$, read $q = \phi'(y)$.

ON THE UNIFORMITY OF THE CONVERGENCE OF CERTAIN ABSOLUTELY CONVERGENT SERIES

BY MAXIME BÔCHER

IF the series

$$(1) \quad u_1(x) + u_2(x) + \dots$$

is absolutely and uniformly convergent for the values of x in a certain interval, and if we rearrange the terms, will the resulting series necessarily be uniformly convergent? This question must be answered in the negative as the following example shows:

$$(2) \quad x^2 - x^2 + \frac{x^2}{1+x^2} - \frac{x^2}{1+x^2} + \frac{x^2}{(1+x^2)^2} - \frac{x^2}{(1+x^2)^2} + \dots$$

Here $S_{2n} = 0$, $S_{2n+1} = \frac{x^2}{(1+x^2)^n}$.

It is easily seen that S_{2n+1} , which reaches its maximum when $x = \pm 1/\sqrt{n-1}$, is always less than $1/n$. Accordingly (2) converges uniformly to the value zero. It is also readily shown to be absolutely convergent.

Now form the series

$$x^2 - x^2 + \frac{x^2}{1+x^2} + \frac{x^2}{(1+x^2)^2} - \frac{x^2}{1+x^2} + \frac{x^2}{(1+x^2)^3} + \frac{x^2}{(1+x^2)^4} - \frac{x^2}{(1+x^2)^5} + \dots$$

from (2) by rearranging the terms. This series, of course, also converges to the value zero, but it converges non-uniformly in any interval including the point $x = 0$. For

$$S_{3n-1} = \frac{x^2}{(1+x^2)^n} \left[1 + \frac{1}{1+x^2} + \dots + \frac{1}{(1+x^2)^{n-2}} \right] = \frac{(1+x^2)^{n-1} - 1}{(1+x^2)^{2n-2}},$$

and when x has either of the real values

$$\pm \sqrt[n-1]{\sqrt{2}-1}$$

we have $S_{3n-1} = 1/4$.

It is, however, easy to prove the following theorem:

If the series

$$|u_1(x)| + |u_2(x)| + \dots$$

is uniformly convergent throughout the interval in question, (1) will be absolutely and uniformly convergent, and will remain so no matter how the order of the terms is changed.

Any series which can be proved uniformly convergent by Weierstrass's test (*cf.* Weierstrass, *Werke*, Vol. II, p. 202) will therefore remain uniformly convergent when its terms are rearranged.

The facts here referred to would seem to be of especial interest in relation to the subject of absolutely convergent multiple series.

In conclusion I will mention the following theorem which I had occasion to prove and to use in a slightly-different form a few years ago (*cf.* *Bull. Amer. Math. Soc.* May, 1898, p. 368):

If throughout the interval $a \leq x \leq b$ the functions $u_i(x)$ are continuous and nowhere negative, and if the function represented by the series (1) is continuous throughout the interval $a \leq x \leq b$, then (1) converges uniformly throughout this interval.